


Concept-Development Practice Page **6-3**

Racing Day with $a = F/m$

In each situation below, Cart A has a mass of **1 kg**. Circle the correct answers (A, B, or Same for both).


1. Cart A is pulled with a force of **1 N**.
 Cart B also has a mass of **1 kg** and is pulled with a force of **2 N**.
 Which undergoes the greater acceleration?

(A) (B) (Same for both)




2. Cart A is pulled with a force of **1 N**.
 Cart B has a mass of **2 kg** and is also pulled with a force of **1 N**.
 Which undergoes the greater acceleration?

(A) (B) (Same for both)




3. Cart A is pulled with a force of **1 N**.
 Cart B has a mass of **2 kg** and is pulled with a force of **2 N**.
 Which undergoes the greater acceleration?

(A) (B) (Same for both)




4. Cart A is pulled with a force of **1 N**.
 Cart B has a mass of **3 kg** and is pulled with a force of **3 N**.
 Which undergoes the greater acceleration?

(A) (B) (Same for both)




5. This time Cart A is pulled with a force of **4 N**.
 Cart B has a mass of **4 kg** and is pulled with a force of **4 N**.
 Which undergoes the greater acceleration?

(A) (B) (Same for both)



6. Cart A is pulled with a force of **2 N**.
 Cart B has a mass of **4 kg** and is pulled with a force of **3 N**.
 Which undergoes the greater acceleration?

(A) (B) (Same for both)



thanx to Dean Baird

© Pearson Education, Inc., or its affiliate(s). All rights reserved.

CONCEPTUAL PHYSICS

Drop and Pull

1. Consider a 1-kg cart being pulled by a 10-N applied force. According to Newton's second law, acceleration of the cart is

$$a = \frac{F}{m} = \frac{10 \text{ N}}{1 \text{ kg}} = 10 \text{ m/s}^2.$$




This is the same as the acceleration of free fall, g —because a force equal to the cart's weight accelerates it.



2. Consider the acceleration of the cart when the applied force is due to a 10-N iron weight attached to a string draped over a pulley. Will the cart accelerate as before, at 10 m/s^2 ? The answer is no, because the mass being accelerated is the mass of the cart *plus* the mass of the piece of iron that pulls it. Both masses accelerate. The mass of the 10-N iron weight is 1 kg—so the total mass being accelerated (cart + iron) is 2 kg. Then,

$$a = \frac{F}{m} = \frac{10 \text{ N}}{2 \text{ kg}} = 5 \text{ m/s}^2.$$

The pulley changes only the direction of the force.



Don't forget; the total mass of a system includes the mass of the hanging iron.




Note this is half the acceleration due to gravity alone, g . So the acceleration of 2 kg produced by the weight of 1 kg is $g/2$.



- a. Find the acceleration of the 1-kg cart when two identical 10-N weights are attached to the string.

$$a = \frac{F}{m} = \frac{\text{applied force}}{\text{total mass}} = \underline{\hspace{2cm}} = \underline{\hspace{2cm}} \text{ m/s}^2$$




Here we simplify and say $g = 10 \text{ m/s}^2$.



CONCEPTUAL PHYSICS

Drop and Pull—continued

- b. Find the acceleration of the 1-kg cart when three identical 10-N weights are attached to the string.




$$a = \frac{F}{m} = \frac{\text{applied force}}{\text{total mass}} = \underline{\hspace{2cm}} = \underline{\hspace{2cm}} \text{ m/s}^2$$

- c. Find the acceleration of the 1-kg cart when four identical 10-N weights (not shown) are attached to the string.

$$a = \frac{F}{m} = \frac{\text{applied force}}{\text{total mass}} = \underline{\hspace{2cm}} = \underline{\hspace{2cm}} \text{ m/s}^2$$

- d. This time 1 kg of iron is added to the cart, and only one iron piece dangles from the pulley. Find the acceleration of the cart.




$$a = \frac{F}{m} = \frac{\text{applied force}}{\text{total mass}} = \underline{\hspace{2cm}} = \underline{\hspace{2cm}} \text{ m/s}^2$$



The force due to gravity on a mass m is mg .
So gravitational force on 1 kg is $(1 \text{ kg})(10 \text{ m/s}^2) = 10 \text{ N}$.

- e. Find the acceleration of the cart when it carries two pieces of iron and only one iron piece dangles from the pulley.




$$a = \frac{F}{m} = \frac{\text{applied force}}{\text{total mass}} = \underline{\hspace{2cm}} = \underline{\hspace{2cm}} \text{ m/s}^2$$

© Pearson Education, Inc., or its affiliate(s). All rights reserved.


- f. Find the acceleration of the cart when it carries 3 pieces of iron and only one iron piece dangles from the pulley.


$$a = \frac{F}{m} = \frac{\text{applied force}}{\text{total mass}} = \underline{\hspace{2cm}} = \underline{\hspace{2cm}} \text{ m/s}^2$$




- g. Find the acceleration of the cart when it carries 3 pieces of iron and 4 pieces of iron dangle from the pulley.

$$a = \frac{F}{m} = \frac{\text{applied force}}{\text{total mass}} = \underline{\hspace{2cm}} = \underline{\hspace{2cm}} \text{ m/s}^2$$



 Mass of cart is 1 kg. Mass of 10-N iron is also 1 kg.



- h. Draw your own combination of masses and find the acceleration. 2 pts

$$a = \frac{F}{m} = \frac{\text{applied force}}{\text{total mass}} = \underline{\hspace{2cm}} = \underline{\hspace{2cm}} \text{ m/s}^2$$

CONCEPTUAL PHYSICS